



Date Palm (*Phoenix dactylifera*): Novel Findings and Future Directions for Food and Drug Discovery



Ali H. El-Far^{1,*}, Babatunji E. Oyinloye², Masood Sepehrimanesh³, Mahmoud A. Gab Allah⁴, Ibrahim Abu-Reidah⁵, Hazem M. Shaheen⁶, Iman Razeghian-Jahromi⁷, Abd el-wahab A. Alsenosy⁸, Ahmed E. Noreldin⁹, Soad K. Al Jaouni¹⁰ and Shaker A. Mousa^{11,*}

¹Department of Biochemistry, Faculty of Veterinary Medicine, Damanhour University, Damanhour 22511, Egypt; ²Phytomedicine, Biochemical Toxicology, and Biotechnology Research Laboratories, Department of Biochemistry, College of Sciences, Afe Babalola University, Ado-Ekiti, 360001 Nigeria; ³Gastrointestinal and Liver Diseases Research Center, Guilan University of Medical Sciences, Rasht, Iran; ⁴Department of Forensic Medicine and Toxicology, Faculty of Veterinary Medicine, University of Sadat City, Sadat, Egypt; ⁵Industrial Chemistry Department, Faculty of Sciences, Arab American University, P.O. Box. 240, 13 Zababdeh-Jenin, Palestine; ⁶Department of Pharmacology, Faculty of Veterinary Medicine, Damanhour University, Damanhour 22511, Egypt; ⁷Cardiovascular Research Center, Shiraz University of Medical Sciences, Shiraz, Iran; ⁸Department of Biochemistry, Faculty of Veterinary Medicine, Damanhour University, Damanhour 22511, Egypt; ⁹Department of Histology and Cytology, Faculty of Veterinary Medicine, Damanhour University, Damanhour, Egypt; ¹⁰Hematology/Pediatric Oncology, King Abdulaziz University Hospital and Scientific Chair of Yousef Abdullatif Jameel of Prophetic Medicine Application, Faculty of Medicine, King Abdulaziz University, Jeddah 21589, Saudi Arabia and ¹¹Pharmaceutical Research Institute, Albany College of Pharmacy and Health Sciences, Rensselaer, NY 12144, USA

ARTICLE HISTORY

Received: October 06, 2017
Revised: January 14, 2018
Accepted: March 15, 2018

DOI:
10.2174/1570163815666180320111937



CrossMark

Abstract: *Phoenix dactylifera* is a useful traditional medicinal plant, mainly the fruit is used, which is the edible part of the plant (Ajwa date). It is now considered to be a valuable source of natural medicinal products against various diseases. Phytochemical investigations have shown that the fruit contains anthocyanins, phenolics, sterols, carotenoids, and flavonoids. The fruits are a rich source of carbohydrates, vitamins, and proteins. *P. dactylifera* is considered as a complete diet because it also contains different fatty acids, amino acids, proteins, and steroidal substances. This review highlights the phytochemical composition, nutritional significance, and potential health benefits of *P. dactylifera* and discusses its potential as a functional food for disease prevention, management, and treatment.

Keywords: *Phoenix dactylifera*, biological activities, phytochemicals, future directions, drug discovery, medicinal plant.

1. INTRODUCTION

Over the years, it has become increasingly evident that nature has numerous medicinal plants that contain phytochemicals with potent pharmacological activities, including promising antitoxic properties [1]. It is estimated by the World Health Organization that up to 80 percent of humans still rely on traditional medicines [2].

P. dactylifera is a useful traditional medicinal plant belonging to the family *Arecaceae*. [3]. The genus *Phoenix* contains 14 species including *P. dactylifera* that have been cultivated in the Middle East for at least 6000 years [4]. Its phytochemical investigation has revealed that the fruits contain anthocyanins, phenolics, sterols, carotenoids,

procyanidins, and flavonoids, compounds known to possess free radical scavenging, antioxidant, antimutagenic, antimicrobial, anti-inflammatory, antihyperlipidemic, gastroprotective, hepatoprotective, nephroprotective, anticancer, and immunostimulant activities [5, 6].

All *P. dactylifera* products such as fruits, seeds, pollen, leaves, and syrup have beneficial uses for human and animals. This attracts researchers to study the different pharmacological potentials of *P. dactylifera* products. Our previous study showed antioxidant and immune stimulant activities of *P. dactylifera* seeds supplemented to broiler chickens as a feed additive [7].

P. dactylifera products are widely used daily food in Islamic countries due to their history of more than 1400 years. Ajwa dates, a special kind of date present in Arab countries, have been mentioned by the Prophet as he said, "If Somebody takes seven Ajwa dates in the morning, neither magic nor poison will hurt him that day" [8]. In addition, the Prophet mentioned the benefits of dates from Medina, a city

*Address correspondence to this author at the Department of Biochemistry, Faculty of Veterinary Medicine, Damanhour University, Damanhour 22511, Egypt; E-mail: ali.elfar@damanhour.edu.eg; Pharmaceutical Research Institute, Albany College of Pharmacy and Health Sciences, Rensselaer, NY 12144, USA; Tel/Fax: 00201009612452; E-mail: shaker.mousa@acphs.edu

in Saudi Arabia, and said, “He who ate seven dates (of the land situated) between these two lava plains in the morning, no poison will harm him until it is evening” [9].

2. *P. DACTYLIFERA* ACTIVE INGREDIENTS

P. dactylifera can be considered as a complete food in addition to its curative effects against various complaints. *P. dactylifera* is considered as a complete diet because it contains different fatty acid, amino acid, proteins, and steroidal substances [10-13]. It contains phytochemicals like phenolics, flavonoids, and carotenoids, which have anticancer and antioxidant potentials [14-16]. The protective effects of *P. dactylifera* are thought to be due to not only the fiber, vitamins, and minerals, but also to a diversity of plant secondary metabolites as flavonoids and phenolics [17, 18]. Also, the polyphenolic proanthocyanidins may act in combination with other phenolics as free radical scavengers or heavy metal chelators, and in turn, they can prevent the oxidative stress and inflammation [19].

P. dactylifera contains considerable amounts of vitamins of water and fat-soluble origins that are very important for vitality. Moreover, it contains vitamins with powerful antioxidant potentials that are capable of chelating different radicals in the non-enzymatic reaction such as vitamin A, C, and E [18, 20]. Additionally, it contains amounts of macro- and micronutrients that play an important role in many biological functions in the body [21-23]. Selenium, copper, zinc, and manganese are of great importance for antioxidant metalloenzymes such as superoxide dismutase (SOD) and glutathione peroxidase (GPx) [24]. The phytochemical constituents of *P. dactylifera* fruits are listed in Tables S1-4 and illustrated in Fig. (1).

3. BIOLOGICAL ACTIVITIES OF *P. DACTYLIFERA*

3.1. Antioxidant Activity

Studies concerning the antioxidant effect of *P. dactylifera* have shown that oxidative stress induced by reactive oxygen species (ROS) and reactive nitrogen species (RNS) reduced the body antioxidant defense system and led to cellular oxidative damage as listed in Table S5. *P. dactylifera* have shown antioxidant activities in the Trolox equivalent antioxidant capacity (TEAC) test, 2,2'-azinobis (3-ethylbenzothiazoline-6-sulphonic acid) radical cation (ABTS⁺) assay, and the ferric reducing/antioxidant power method (FRAP assay) [25]. Also, *P. dactylifera* contains a high percentage of vitamin C, A, and E plus a high total phenolic content [26, 27]. These results of *in vitro* studies encouraged researchers to investigate the antioxidant activity of *P. dactylifera* extracts *in vivo* against different toxicants as carbon tetrachloride (CCl₄), isoproterenol, cadmium, and streptozotocin-induced diabetic rats [28-31]. The protective effect of *P. dactylifera* may be related to the accelerated activities of antioxidant enzymes such as catalase (CAT), SOD, GPx, glutathione reductase (GR), glutathione S-transferase (GST) along with significant reduction in malondialdehyde (MDA) [7, 30, 32, 33]. Conclusively, it can be said that *P. dactylifera* is a good antioxidant food.

3.2. Anticancer Activity

Constituents of *P. dactylifera* fruits have shown antitumor activity, as listed in Table S6. *In vitro* trials were done to determine the anticancer activity of *P. dactylifera* extracts towards different cancer cell lines such as human epithelial colorectal adenocarcinoma (Caco-2) [34], and the human melanoma-derived cell line (IGR-39) [35]. Khan and colleagues investigated the mechanism by which the methanolic extract of Ajwa date (15 and 20 mg/ml) inhibited the growth of human breast adenocarcinoma (MCF7) through upregulation of proapoptotic molecules, p53, Bcl-2-associated X protein (Bax), Fas, and Fas ligand (FasL) along with downregulation of B-cell lymphoma 2 (Bcl-2) [36].

3.3. Antidiabetic, Anti-Hyperglycemic, and Antihyperlipidemic Activities

We found four studies in the literature on the antidiabetic activity of *P. dactylifera* (Table S7). Some studies proofed the antidiabetic effect of *P. dactylifera* through elevation of plasma insulin with normalization of plasma glucose, triacylglycerol, and cholesterol in alloxan-induced diabetes in rats [37]. It is important to investigate the real cause of plasma insulin increment as to whether this is due to increase in the efficiency of healthy β -cells or due to the regeneration of alloxan-injured cells.

One of the antihyperglycemic effects of *P. dactylifera* is the inhibition of either α -glucosidase and α -amylase *in vitro*, potentially delaying the digestion and absorption of carbohydrates, leading to normalization of plasma glucose levels. Such an effect was induced by *P. dactylifera* leaves' hydro-alcoholic extract [35] and *P. dactylifera* seeds' aqueous extract [38]. Also, Hasan and Mohieldein [39] treated streptozotocin-induced diabetic rats with 10 ml of *P. dactylifera* seeds' aqueous extract per day for each rat. The tested extract significantly reverted the elevated serum glucose, cholesterol, and triacylglycerol levels of the diabetic rats to near normal values.

3.4. Anti-Inflammatory Activity

The anti-inflammatory effects of *P. dactylifera* could be shown as a reduction of paw edema volume [40], wound healing activities [41], and cardioprotective [30]. In line with these facts, we summarize the anti-inflammatory properties of *P. dactylifera* as reported in the literature (Table S8).

Al-Qarawi and coworkers found that aqueous and ethanolic undialyzed and dialyzed extracts from *P. dactylifera* fruit have been effective in ameliorating the severity of gastric ulceration and mitigating the ethanol-induced increase in histamine and gastrin concentrations and decreasing mucin gastric levels. According to the authors, the anti-inflammatory ability of the extract may be explained by the antioxidant action of the extract [42].

The phenolics and flavonoids content could inhibit the formation of prostaglandin endoperoxide, leading to termination of inflammation mediators like prostaglandins and thromboxane [43]. Ali Haimoud and colleagues evaluated the anti-inflammatory activity of methanolic extracts of




Fig. (1). The structures of some secondary metabolites characterized in *P. dactylifera*.

P. dactylifera fruits grown in Algeria. They found a noticeable reduction in paw volume in the experimental model of carrageenan-induced acute paw edema in Swiss albino mice ranging from 35.64 to 67.56% as a response to oral administration of 250 mg/kg dose of the methanolic extract [40].

In traditional medicine, *P. dactylifera* has been used to treat inflammatory-associated diseases [44] and recently was presented as a pain reliever along with chemical agents such as ibuprofen and paracetamol [45, 46]. There is no clear mechanism for such effects but it has been reported that *P. dactylifera* had active substances that can interfere with prostaglandin synthesis [47], inhibiting the expression of inflammatory cytokines such as IL-6, IL-8, IL-10, TNF- α , and IGF-1 [30], analgesic effects due to the presence of vitamin C and E [46], and increasing the expression of TGF- β [48]. On the other hand, potentiating of the antioxidant

system as described previously can be a possible mechanism for anti-inflammatory effects of *P. dactylifera*.

3.5. Antimicrobial Activity

Different extracts and oils of *P. dactylifera* show strong antimicrobial activities (Table S9).

Abuharfeil *et al.* [49] studied the effect of *P. dactylifera* against the hemolytic activity of streptolysin O and stated that the *P. dactylifera* fruit extract resulted in a decrease in the growth of *Streptococcus pyogenes* by 88.5% compared to control. In a disc diffusion method, it has been determined that aqueous and ethanol extracts of *P. dactylifera* fruits had a strong antibacterial activity against *Escherichia coli*, *Salmonella enterica*, and *Bacillus subtilis* and moderately inhibited *Staphylococcus aureus* and *Enterococcus faecalis*. This antibacterial effect is due to the presence of esculetin,

tannic acid, a moderate concentration of gallic acid, itaconic acid, and traces of ferulic acid present in *P. dactylifera* extracts [50].

In another study, the antibacterial activity of methanol and acetone extract of the Saudi Arabian *P. dactylifera* variety Mabroom, Safawi, and Ajwa, as well as Iranian *P. dactylifera* variety Mariami were evaluated. The methanolic extract of Mabroom was more potent against *Staphylococcus aureus* than the methanolic extract of Ajwa and Mariami (100 mg/ml for Mabroom vs. 300 mg/ml for Ajwa and Mariami). The lowest concentration of methanolic extracts of Mabroom and Mariami that could inhibit *Bacillus cereus* growth was 500 mg/ml while methanolic extracts of Safawi and Ajwa could inhibit the growth of this bacterial species at 400 mg/ml [27].

Finally, antimicrobial activities of hydroalcoholic extract of six Moroccan date fruit varieties including Bouskri, Bousrdon, Boushammi, Boufgous, Jihl, and Majhoul against Gram-positive (*Bacillus subtilis*, *B. cereus*, and *S. aureus*) and Gram-negative bacteria (*E. coli*, *Pseudomonas aeruginosa*, *Salmonella abony*) were examined by Bouhlali and colleagues using a disc diffusion method. All the tested *P. dactylifera* fruit extracts showed antibacterial activity except Majhoul and Bouskri extracts, but this antibacterial effect was still lower than that of gentamicin. The Bousrdon and Jihl extracts were more potent inhibitors with minimum inhibitory concentration (MIC) values ranging from 2.5-10 mg/ml for all bacterial strains tested [51].

As noted above, *P. dactylifera* can be effective against Gram-positive and -negative bacteria. However, it has been reported that *P. dactylifera* fruits are more efficient against Gram-positive bacteria than against Gram-negative bacteria due to the presence of an outer membrane. On the other hand, the most effective substances in *P. dactylifera* that are attributed to the antimicrobial effects are phenolic compounds by generating hydrogen peroxide that mediates bacterial growth inhibition [52]. It has been postulated that phenolic compounds utilize redox active metals when interacting with bacteria, in particular, gram-positive [47].

3.6. Antitoxic Activities

The studies concerning the hepatoprotective, nephroprotective, and neuroprotective effects of *P. dactylifera* against oxidative stress, and xenobiotics-induced toxicity are listed in Table S10.

3.6.1. Antihepatotoxicity

The hepatoprotective potential of *P. dactylifera* was examined in a study by Al-Qarawi and coworkers on the effect of aqueous extracts of the flesh and seeds against CCl₄-induced hepatotoxicity in rats. The *P. dactylifera* seeds' extract was added to drinking water. β -Sitosterol, one of the bioactive compounds present in the extract, may be responsible for the observed protective effect against CCl₄-induced hepatic injury in a rat model [53]. In screening and developing of hepatoprotective drugs for the treatment of hepatocellular injuries, the ability of the constituents of such drugs to inhibit the aromatase activity of cytochrome P-450, thereby favoring liver regeneration, is an important factor to consider [54]. Based on the aforementioned results,

Kowalska *et al.* in 1990 suggested that the flavonoids content of *P. dactylifera* could be a factor contributing to its hepatoprotective ability through the inhibition of cytochrome P-450 aromatase [55].

Saafi and coworkers reported a protective effect of the aqueous fruit flesh extract of *P. dactylifera* against dimethoate-induced oxidative stress in rat liver. Oral administration of dimethoate (20 mg/kg) resulted in the elevation of all hepatic biomarker enzymes examined. The elevation witnessed in the hepatic biomarker enzymes examined was an indication of a certain degree of damage to the liver. Taken together, their results demonstrated that when animals challenged with dimethoate are treated with aqueous fruit flesh extract (4 ml/kg) of *P. dactylifera*, the extract can mitigate diminished hepatic antioxidant activities, notably SOD, GPx, and CAT activities, and inhibit hepatic lipid peroxidation as well as restore various degrees of alterations observed in hepatic biomarker enzymes, thereby preventing against hepatocellular damage [56].

3.6.2. Antinephrotoxicity

P. dactylifera has also been reported to play a crucial role in the treatment and management of various nephrotoxicities arising from different xenobiotics. In a study conducted by Ahmed and coworkers, using a CCl₄-induced toxicity model in rats, a hydroacetone extract (50 or 100 mg/kg/rat) prepared from seeds of *P. dactylifera* was demonstrated to confer a noticeable protection on the kidney in a dose-dependent manner. According to the authors, the nephroprotective ability of the extract may be explained by its ability to effectively scavenge the free radical generated during CCl₄ metabolism, probably because the extract contained a high quantity of proanthocyanidins (49.0 % CT – 0.52 % CT w/w), which exhibited high antioxidant activity [57]. Another study conducted by Hasan and Mohieldin revealed that prolonged administration of aqueous *P. dactylifera* extracts (100 g/L in a dosage of 10 ml/day/rat) could restore kidney function. In this study, it was observed that there were various degrees of renal dysfunction among diabetic rats while treatment with aqueous *P. dactylifera* extracts ameliorated the progressive decline in renal dysfunction among the treated rats [39].

Al-Ghasham and coworkers reported that aflatoxin B₁ (AFB₁) induces histopathological changes in the kidney that are associated with renal dysfunction, as indicated by the elevations in the levels of plasma creatinine and urea. Toxicity was induced by intraperitoneal administration of AFB₁ (50 μ g/kg). Treatment with aqueous *P. dactylifera* extract for two weeks resulted in a marked reduction in levels of plasma creatinine and urea as well as a significant improvement in the kidney architecture. They concluded that the observed effect may be attributed to the antioxidant properties of *P. dactylifera* extract [58]. Al-Qarawi and coworkers also investigated the effect of *P. dactylifera* extract on gentamicin (GM) nephrotoxicity in rats. Toxicity was induced by administration of GM (80 mg/kg/day) intramuscularly. *P. dactylifera* extracts were administered to the animals by mixing the *P. dactylifera* flesh extract with the food (50 % w/w) or by mixing the seed extract in the drinking water (2:1 w/v). GM administration was found to cause a significant elevation in the plasma levels of

creatinine and urea and induced a marked necrosis of the renal proximal tubules. Interestingly, administration of either *P. dactylifera* flesh extracts with the food (50% w/w) or seeds extract in the drinking water (2:1 w/v) resulted in a significant reversal effect in observed indices of toxicity in the kidney. Melatonin, vitamin E, and ascorbic acid, which are abundantly present in the extract and may synergistically act to counteract the overwhelming effect of the free radicals generated, were suggested to be the basis of the nephroprotection [59].

3.6.3. Antineurotoxicity

The impact of free radical generation, mediated by oxidative stress in the pathogenesis of various neurological diseases such as a cognitive decline in aging, Alzheimer's disease, Parkinson's disease, and vascular dementia has been widely documented in the literature [60]. Antioxidants from plant-derived food in the form of nutraceuticals are now being considered as an alternative therapy against solemn neuronal loss due to their capability to counteract free radicals generated and neutralize them, thereby conferring neuroprotection [61]. Pujari and co-workers investigated the neuroprotective and antioxidant effect of *P. dactylifera* fruits against permanent bilateral common carotid arteries (BCCA) in rats at doses of 30, 100, and 300 mg/kg. They reported that chronic occlusion of BCCA resulted in a noticeable increase in lipid peroxidation as evidenced by the elevation in malondialdehyde levels. They also observed that there was a general decline in the endogenous antioxidants, notably glutathione, GPx, GR, GST, CAT, and SOD. They concluded that 100 and 300 mg/kg of the extract significantly ameliorated these alterations, confirming the protective role of the extract in ischemia hypoperfusion [62]. The polyphenolic constituents in the extract such as flavonoids and plant sterols as well as its ascorbic acid content may account for the observed neuroprotective effect.

A similar study conducted to investigate the neuroprotective effect of aqueous *P. dactylifera* fruit extract in focal cerebral ischemia in rats reported that a 250 mg/kg dose of the extract significantly inhibited neuronal damage induced by cerebral ischemia. Rats were maintained on varying doses of the extract (125, 250, 500, and 1000 mg/kg). The extract was administered once per day for two weeks; the largest protective effects of the extract were observed at a dose of 250 mg/kg. At 500 mg/kg, a lower protective effect was observed, and 1000 mg/kg showed a negative effect, which may be attributed to high concentration of antioxidants that may be harmful. They concluded that the extract at 250 mg/kg could protect neurons against ischemia-reperfusion-induced insults [63]. Taken together, in search for a promising antitoxic agent in the treatment and management of hepatotoxicity, nephrotoxicity, and neurotoxicity, *P. dactylifera* is a candidate to explore.

3.7. Reproductive and Labor Induction Activities

Plants and their constituents play a significant role as pain relievers and cause relaxation in childbirth. Several medicinal plants show an effect in the stimulation of all phases of labor [64]. *P. dactylifera* has a pivotal role in the

enhancement of male fertility and induction of labor as stated in Table S11 that may be related to its antioxidant activity [65, 66].

An *in vivo* trial was started to evaluate the effect of the normal and acid-treated powdered *P. dactylifera* seeds on male rat fertility [67]. The normal *P. dactylifera* seeds concentrations of 7 and 14% caused a significant increase in the plasma testosterone, while the acid-treated seeds (14%) significantly increased the plasma level of luteinizing hormone (LH). In another study, the authors injected 0.3 ml of diluted *P. dactylifera* seeds oil (15 and 20% of oil in saline 0.9%) and reported significant increases in the sperm count, motility, and viability in male mice [68].

Another study reported the reproductive potential of *P. dactylifera* pollen through treatment of rats by 1 ml of aqueous extract of pollen using concentrations of 120, 240, 360 mg/kg body weight by gavage. The doses of 120 and 240 mg/kg significantly raised the ratio of testis or epididymis to body weight, sperm count, sperm motility, and estradiol level compared to the control group, with noticeable increases in serum LH and testosterone levels [69]. Where LH and testosterone are responsible for spermatogenesis and male fertility, estradiol that is produced by Leydig cells of the mammalian testis prevents the apoptosis of male sperm cells. These findings were confirmed by the study of El-Kashlan, *et al.* [70] in which they administered 150 mg ethanolic extract of *P. dactylifera* pollen per kg in male rat and recorded marked an increase in sperm count and motility, serum levels of LH, testosterone, and estradiol hormones. The authors also recognized potentiation in the activities of testicular 3β -hydroxysteroid dehydrogenase (3β -HSD) and 17β -hydroxysteroid dehydrogenase (17β -HSD) that contribute to the testosterone biosynthesis.

3.8. Cardiovascular Protective Activity

P. dactylifera fruit has been used as an antihypertensive food for centuries. Because *P. dactylifera* seeds can be used as edible oils and also in pharmaceutical industry, it has been proposed that it may exert beneficial effects on cardiovascular conditions [20]. Braga *et al.* showed that this fruit is a potent angiotensin-converting enzyme inhibitor. This is an effective strategy for reducing blood pressure [71]. Also, high sodium and low potassium intake have a major role in raising blood pressure [72]. Because *P. dactylifera* fruit contains a high amount of potassium and a low amount of sodium, it can control blood pressure by maintaining the electrolyte balance. Also, the magnesium and calcium contents of *P. dactylifera* fruit play key roles in this subject [73].

Hypercholesterolemia is an important risk factor because of its major impact on the progression of cardio- and cerebrovascular disorders. It was reported that *P. dactylifera* fruit supplementation could modulate cholesterol absorption and metabolism [74]. This effect is related to *P. dactylifera* fiber and phytochemicals by three mechanisms: reduction in cholesterol absorption and reabsorption of bile acids, and inhibition of hepatic cholesterol biosynthesis after production of short-chain fatty acids due to *P. dactylifera* fiber fermentation. Also, phytochemicals such as phytosterols and

phytoestrogens inhibit cholesterol absorption from intestinal lumen [75].

Ahmed *et al.* [76] demonstrated that for *Aseel*, the best varieties of Pakistani dates, efficiency was the same as atorvastatin with respect to lowering of some markers such as fasting blood sugar, cholesterol, triacylglycerol, LDL, and VLDL. Generally, this experiment showed that *Aseel* lowered the blood lipid levels. The presence of some constituents like phytochemicals, β -sitosterol, proanthocyanidin, catechin, quercetin, anthocyanins, and selenium may contribute to cardioprotective and antihyperlipidemic effects of *P. dactylifera* [77].

4. FUTURE DIRECTIONS

Suggestions for future research directions to give a complete understanding of the biological and nutritive roles of *P. dactylifera* are:

- Phytochemical constituents of *P. dactylifera* grown in Arabian countries.
- Comparative phytoconstituents of Aliya, Medina and other countries (The Middle East and North Africa).
- Differences between *P. dactylifera* eaten in the early morning (empty stomach) and other times of the day.
- Protective effect of *P. dactylifera* against different toxicants in different models.
- Plasma amino acid pool, growth hormone, and insulin-like growth hormone levels.
- Growth hormone receptor and myogenic genes in muscles.
- Plasma lipid profiles and non-esterified fatty acids (NEFA).
- Experimentally induced non-alcoholic fatty liver concerning triacylglycerol and cholesterol biosynthesis-related genes.
- The relationship between *P. dactylifera* and intestinal amino acid transporters such as oligopeptide transporter (PepT1), excitatory amino acid transporter 3 (EAAT3), Na^+ -independent branched-chain and aromatic amino acid transporter (LAT1), Na^+ -independent cationic amino acid transporter (CAT1), Na^+ -independent cationic amino acid transporter (CAT2).
- Purification of active substances responsible for antioxidant activity.
- Application of different extracts of *P. dactylifera* in different routes as an alternative treatment for different diseases with a base of oxidative stress.
- Pro-apoptotic proteins such as caspase-3, caspase-8, caspase-9, phosphatase and tensin homolog (PTEN), p53 upregulated modulator of apoptosis (PUMA), phorbol-12-myristate-13-acetate-induced protein 1 (NOXA), integrins, and E-cadherin.
- Anti-apoptotic molecules like mitogen-activated protein kinase (MAPK), nuclear factor kappa B (NF- κ B), actin, cyclin-dependent kinases (CDKs), matrix metalloproteinases (MMPs), mammalian target of rapamycin (mTOR), signal transducer and activator of transcription 3 (STAT3), and receptor for advanced glycation end-products (RAGE) signaling pathways.
- The combination between *P. dactylifera* and the already established anticancer drugs of either natural or synthetic nature.
- The relationship between *P. dactylifera* and intestinal monosaccharide transporters such as Na^+ -dependent glucose and galactose transporter (SGLT1), glucose transporter (SGLT5), Na^+ -independent glucose, galactose and fructose transporter (GLUT2), and Na^+ -independent fructose transporter (GLUT5).
- Glucose uptake by peripheral tissues with evaluation of glucokinase and hexokinase genes and protein expression.
- Gluconeogenic, glycogenic, and glycogenolytic enzymes' gene and protein expression.
- Fatty acid β -oxidation enzymes' gene and protein expression.
- Regeneration of injured β -cells in diabetic-induced models.
- Beneficial effects of extracts of *P. dactylifera* in the experimental model of human inflammatory diseases.
- Proteomic analysis of differentially expressed proteins in the liver, kidney, and brain of animal models in xenobiotics-induced toxicity (CCl_4 or ethanol) should be investigated as identification of these differentially expressed proteins will help us to understand the molecular events occurring during xenobiotics' toxicity and will also help us in elucidating the exact mechanism of action of *P. dactylifera*.
- Molecular mechanisms of *P. dactylifera* reproductive activity through evaluation of the gene expression of the enzymes of testosterone and estradiol biosynthesis.
- Effect of *P. dactylifera* on female reproductive enzymes such as LH, follicle-stimulating hormone (FSH), estrogen, and progesterone.

- The role of *P. dactylifera* in the regulation of irregular menstrual cycle and silent heat in human and animals.
- The role of *P. dactylifera* in labor induction concerning oxytocin, gonadotropin-releasing hormone (GnRH), and Prostaglandin F₂α (PGF₂α).

CONSENT FOR PUBLICATION

Not applicable.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

Declared none.

SUPPLEMENTARY MATERIAL

Supplementary material is available on the publisher's web site along with the published article.

REFERENCES

- [1] Ekor M. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. *Front Pharmacol* 2014; 4: 177.
- [2] Yadav K, Kadam P, Patel J, Patil M. *Strychnos potatorum*: Phytochemical and pharmacological review. *Pharmacogn Rev* 2014; 8(15): 61-6.
- [3] Sirisena S, Ng K, Ajlouni S. The emerging Australian date palm industry: Date fruit nutritional and bioactive compounds and valuable processing by-products. *Compr Rev Food Sci Food Saf* 2015; 14(6): 813-23.
- [4] Copley MS, Rose PJ, Clapham A, Edwards DN, Horton MC, Evershed RP. Detection of palm fruit lipids in archaeological pottery from Qasr Ibrim, Egyptian Nubia. *Proc Biol Sci* 2001; 268(1467): 593-7.
- [5] Baliga MS, Baliga BRV, Kandathil SM, Bhat HP, Vayalil PK. A review of the chemistry and pharmacology of the date fruits (*Phoenix dactylifera* L.). *Food Res Int* 2011; 44(7): 1812-22.
- [6] El-Far AH, Shaheen HM, Abdel-Daim MM, Al Jaouni SK, Mousa SA. Date palm (*Phoenix dactylifera*): Protection and remedy food. *Curr Tren in Nutra* 2016; 1 (29): 1-10.
- [7] El-Far AH, Ahmed HA, Shaheen HM. Dietary supplementation of *Phoenix dactylifera* seeds enhances performance, immune response, and antioxidant status in broilers. *Oxid Med Cell Longev* 2016; 2016: 9.
- [8] Al-Bukhari MI. *The collection of authentic sayings of Prophet Mohammad (peace be upon him), division 71 on medicine*. In: *Al-Bukhari S. Hilal Yayinlari*, Ankara: Turkey, 1976.
- [9] Siddiqui A. *Sahih Muslim, Book 23: The Book of Drinks, Chapter 25: Excellence of the dates of Medina*. Translator: *Abd-al-Hamid Siddiqui*, 2009.
- [10] Boukouada M, Yousfi M. Phytochemical study of date seeds lipids of three fruits (*Phoenix dactylifera* L) produced in Ouargla region. *Annales de la Faculté des Sciences et Sciences de l'Ingénieur* 2009; 1: 66-74.
- [11] Assirey EAR. Nutritional composition of fruit of 10 date palm (*Phoenix dactylifera* L.) cultivars grown in Saudi Arabia. *J Taibah Univ Med Sci* 2015; 9 (1): 75-9.
- [12] Ali HS, Alhaj OA, Al-Khalifa AS, Bruckner H. Determination and stereochemistry of proteinogenic and non-proteinogenic amino acids in Saudi Arabian date fruits. *Amino Acids* 2014; 46(9): 2241-57.
- [13] Boudries H, Kefalas P, Hornero-Mendez D. Carotenoid composition of Algerian date varieties (*Phoenix dactylifera*) at different edible maturation stages. *Food Chem* 2007; 101 (4): 1372-7.
- [14] Kchaou W, Abbès F, Attia H, Besbes S. *In vitro* antioxidant activities of three selected dates from tunisia (*Phoenix dactylifera* L.). *J Chem* 2014; 2014: 1-8.
- [15] Al-Rimawi F, Odeh I. Development and validation of an HPLC-UV method for determination of eight phenolic compounds in date palms. *J AOAC Int* 2015; 98(5): 1335-9.
- [16] Al-Farsi M, Alasalvar C, Morris A, Baron M, Shahidi F. Comparison of antioxidant activity, anthocyanins, carotenoids, and phenolics of three native fresh and sun-dried date (*Phoenix dactylifera* L.) varieties grown in Oman. *J Agric Food Chem* 2005; 53(19): 7592-9.
- [17] Benmeddour Z, Mehinagic E, Le Meurlay D, Louaileche H. Phenolic composition and antioxidant capacities of ten Algerian date (*Phoenix dactylifera* L.) cultivars: A comparative study. *J Funct Foods* 2013; 5(1): 346-54.
- [18] Vayalil PK. Date fruits (*Phoenix dactylifera* Linn): An emerging medicinal food. *Crit Rev Food Sci Nutr* 2012; 52(3): 249-71.
- [19] Blade C, Aragonés G, Arola-Arnal A, Muguera B, Bravo FI, Salvado MJ, Arola L, Suarez M. Proanthocyanidins in health and disease. *Biofactors* 2016; 42(1): 5-12.
- [20] Al-Shahib W, Marshall RJ. The fruit of the date palm: Its possible use as the best food for the future? *Int J Food Sci Nutr* 2003; 54(4): 247-59.
- [21] Ahmed I. Chemical composition of date varieties as influenced by the stage of ripening. *Food Chem* 1995; 54(3): 305-9.
- [22] Ismail B, Haffar I, Baalbaki R, Mechref Y, Henry J. Physico-chemical characteristics and total quality of five date varieties grown in the United Arab Emirates. *Int J Food Sci Tech* 2006; 41(8): 919-26.
- [23] AlHooti S, Sidhu JS, Qabazard H. Studies on the physicochemical characteristics of date fruits of five UAE cultivars at different stages of maturity. *Arab Gulf J Sci Res* 1995; 13(3): 553-69.
- [24] Adelekan DA, Thurnham DI. Glutathione peroxidase (EC 1.11.1.9) and superoxide dismutase (EC 1.15.1.1) activities in riboflavin-deficient rats infected with *Plasmodium berghei* malaria. *Br J Nutr* 1998; 79(3): 305-9.
- [25] Biglari F, AlKarkhi AFM, Easa AM. Antioxidant activity and phenolic content of various date palm (*Phoenix dactylifera*) fruits from Iran. *Food Chem* 2008; 107(4): 1636-41.
- [26] Hasan NS, Amom ZH, Nor AI, Mokhtarrud N, Mohd Esa N, Azlan A. Nutritional composition and *in vitro* evaluation of the antioxidant properties of various dates extracts (*Phoenix Dactylifera* L.) from libya. *Asia J Clin Nutr* 2010; 2(4): 208-14.
- [27] Samad M, Hashim S, Simarani K, Yaacob J. Antibacterial properties and effects of fruit chilling and extract storage on antioxidant activity, total phenolic and anthocyanin content of four date palm (*Phoenix dactylifera*) cultivars. *Molecules* 2016; 21(4): 419.
- [28] Abdelaziz DH, Ali SA. The protective effect of *Phoenix dactylifera* L. seeds against CCl₄-induced hepatotoxicity in rats. *J Ethnopharmacol* 2014; 155(1): 736-43.
- [29] Abdelaziz DH, Ali SA, Mostafa MM. *Phoenix dactylifera* seeds ameliorate early diabetic complications in streptozotocin-induced diabetic rats. *Pharm Biol* 2015; 53(6): 792-9.
- [30] Al-Yahya M, Raish M, AlSaid MS, Ahmad A, Mothana RA, Al-Sohaibani M, Al-Dosari MS, Parvez MK, Rafatullah S. 'Ajwa' dates (*Phoenix dactylifera* L.) extract ameliorates isoproterenol-induced cardiomyopathy through downregulation of oxidative, inflammatory and apoptotic molecules in rodent model. *Phytomedicine* 2016; 23(11): 1240-8.
- [31] El-Neweshy MS, El-Maddawy ZK, El-Sayed YS. Therapeutic effects of date palm (*Phoenix dactylifera* L.) pollen extract on cadmium-induced testicular toxicity. *Andrologia* 2013; 45(6): 369-78.
- [32] Hoseinifar SH, Dadar M, Khalili M, Cerezuela R, Esteban MÁ. Effect of dietary supplementation of palm fruit extracts on the transcriptomes of growth, antioxidant enzyme and immune-related genes in common carp (*Cyprinus carpio*) fingerlings. *Aquacult Res* 2017; 48(7): 3684-92.
- [33] Sharifi M, Bashtani M, Naserian AA, Farhangfar H. The Effect of increasing levels of date palm (*Phoenix dactylifera* L.) seed on the performance, ruminal fermentation, antioxidant status and milk fatty acid profile of Saanen dairy goats. *J Anim Physiol Anim Nutr* 2017; 101(5): e332-e341.

- [34] Eid N, Enani S, Walton G, Corona G, Costabile A, Gibson G, Rowland I, Spencer JPE. The impact of date palm fruits and their component polyphenols, on gut microbial ecology, bacterial metabolites and colon cancer cell proliferation. *J Nutr Sci* 2014; 3: e46.
- [35] Chakroun M, Khemakhem B, Mabrouk HB, *et al.* Evaluation of anti-diabetic and anti-tumoral activities of bioactive compounds from *Phoenix dactylifera* L's leaf: *in vitro* and *in vivo* approach. *Biomed Pharmacother* 2016; 84: 415-22.
- [36] Khan F, Ahmed F, Pushparaj PN, Abuzenadah A, Kumosani T, Barbour E, AlQahtani M, Gauthaman K. Ajwa Date (*Phoenix dactylifera* L.) Extract inhibits human breast adenocarcinoma (MCF7) cells *in vitro* by inducing apoptosis and cell cycle arrest. *PLoS One* 2016; 11(7): e0158963.
- [37] Mard SA, Jalalvand K, Jafarnejad M, Balochi H, Naseri MK. Evaluation of the antidiabetic and antilipaemic activities of the hydroalcoholic extract of *Phoenix dactylifera* palm leaves and its fractions in alloxan-induced diabetic rats. *Malays J Med Sci* 2010; 17(4): 4-13.
- [38] Khan SA, Al Kiyumi AR, Al Sheidi *et al.* *In vitro* inhibitory effects on α -glucosidase and α -amylase level and antioxidant potential of seeds of *Phoenix dactylifera* L. *Asian Pac J Trop Biomed* 2016; 6(4): 322-9.
- [39] Hasan M, Mohieldein A. *In Vivo* Evaluation of anti diabetic, hypolipidemic, antioxidative activities of saudi date seed extract on streptozotocin induced diabetic rats. *J Clin Diagn Res* 2016; 10(3): FF06-12.
- [40] Ali Haimoud S, Allem R, Merouane A. Antioxidant and anti-inflammatory properties of widely consumed date palm (*Phoenix dactylifera* L.) fruit varieties in Algerian oases. *J Food Biochem* 2016; 40(4): 463-71.
- [41] Abdennabi R, Bardaa S, Mehdi M, *et al.* *Phoenix dactylifera* L. sap enhances wound healing in Wistar rats: Phytochemical and histological assessment. *Int J Biol Macromol* 2016; 88: 443-50.
- [42] Al-Qarawi AA, Abdel-Rahman H, Ali BH, Mousa HM, El-Mougy SA. The ameliorative effect of dates (*Phoenix dactylifera* L.) on ethanol-induced gastric ulcer in rats. *J Ethnopharmacol* 2005; 98(3): 313-7.
- [43] Zhang CR, Aldosari SA, Vidyasagar PS, Nair KM, Nair MG. Antioxidant and anti-inflammatory assays confirm bioactive compounds in Ajwa date fruit. *J Agric Food Chem* 2013; 61(24): 5834-40.
- [44] Yasin B, El-Fawal H, Mousa S. Date (*Phoenix dactylifera*) polyphenolics and other bioactive compounds: A traditional islamic remedy's potential in prevention of cell damage, cancer therapeutics and beyond. *Int J Mol Sci* 2015; 16(12): 30075-90.
- [45] Sani I, Bakar N, Rohin M, Suleiman I, Umar M, Mohamad N. *Phoenix dactylifera* linn as a potential novel anti-oxidant in treating major opioid toxicity. *J Appl Pharm Sci* 2015; 167-72.
- [46] Maryam UI, Simbak N, Umar A, Sani IH, Baig AA, Zin T, Swethadri GKM. Anti-inflammatory and analgesic activities of aqueous extract date palm (*Phoenix dactylifera* L) fruit in rats. *International journal of novel research in healthcare and nursing* 2015; 2: 166-72.
- [47] Taleb H, Maddocks SE, Morris RK, Kanekanian AD. Chemical characterisation and the anti-inflammatory, anti-angiogenic and antibacterial properties of date fruit (*Phoenix dactylifera* L.). *J Ethnopharmacol* 2016; 194: 457-68.
- [48] Elberry AA, Mufti ST, Al-Maghrabi JA, *et al.* Anti-inflammatory and antiproliferative activities of date palm pollen (*Phoenix dactylifera*) on experimentally-induced atypical prostatic hyperplasia in rats. *J Inflamm* 2011; 8(1): 40.
- [49] Abuharfeil NM, Sukhon SE, Msameh Y, Sallal A-KJ. Effect of Date Fruits, *Phoenix Dactylifera* L., on the Hemolytic Activity of Streptolysin O. *Pharm. Biol* 2008; 37(5): 335-9.
- [50] El Sohaimy S, Abdelwahab A, Brennan C, Aboul-enein A. Phenolic content, antioxidant and antimicrobial activities of Egyptian date palm (*Phoenix dactylifera* L.) fruits. *Aust J Basic Appl Sci* 2015; 9(1): 141-7.
- [51] Bouhlali ET, Bammou M, Sellam K, Benlyas M, Alem C, Filali-Zegzouti Y. Evaluation of antioxidant, antihemolytic and antibacterial potential of six Moroccan date fruit (*Phoenix dactylifera* L.) varieties. *J King Saud Univ - Sci* 2016; 28(2): 136-42.
- [52] Taleb H, Maddocks SE, Morris RK, Kanekanian AD. The antibacterial activity of date syrup polyphenols against *S. aureus* and *E. coli*. *Front Microbiol* 2016; 7: 198.
- [53] Al-Qarawi AA, Mousa HM, Ali BH, Abdel-Rahman H, El-Mougy SA. Protective effect of extracts from dates (*Phoenix dactylifera* L.) on carbon tetrachloride-induced hepatotoxicity in rats. *Int J Appl Res Vet Med* 2004; 2(3): 176-80.
- [54] Bastway AM, Hasona NA, Selemain AH. Protective effects of extract from dates (*Phoenix dactylifera* L.) and ascorbic acid on thioacetamide-induced hepatotoxicity in rats. *Iran J Pharm Res* 2008; 7(3): 193-201.
- [55] Kowalska DP, Feeley RM, Jones DP. Use of exogenous glutathione for metabolism of peroxidized methyl linoleate in rat small intestine. *J Nutr* 1990; 120: 1115-21.
- [56] Saafi EB, Louedi M, Elfeki A, Zakhama A, Najjar MF, Hammami M, Achour L. Protective effect of date palm fruit extract (*Phoenix dactylifera* L.) on dimethoate induced-oxidative stress in rat liver. *Exp Toxicol Pathol* 2011; 63(5): 433-41.
- [57] Ahmed AF, Al-Qahtani JH, Al-Yousef HM, Al-Said MS, Ashour AE, Al-Sohaibani M, Rafatullah S. Proanthocyanidin-rich date seed extract protects against chemically induced hepatorenal toxicity. *J Med Food* 2015; 18(3): 280-9.
- [58] Al-Ghsham A, Ata HS, El-Deep S, Meki AR, Shehada S. Study of protective effect of date and *Nigella sativa* on aflatoxin b(1) toxicity. *Int J Health Sci (Qassim)* 2008; 2(2): 26-44.
- [59] Al-Qarawi AA, Abdel-Rahman H, Mousa HM, Ali BH, El-Mougy SA. Nephroprotective Action of *Phoenix dactylifera*. in Gentamicin-Induced Nephrotoxicity. *Pharm. Biol* 2008; 46(4): 227-30.
- [60] Uttara, B.; Singh, A.; Zamboni, P.; Mahajan, R., Oxidative Stress and Neurodegenerative Diseases: A Review of Upstream and Downstream Antioxidant Therapeutic Options. *Curr Neuropharmacol* 2009; 7(1): 65-74.
- [61] Pujari RR, Vyawahare NS, Kagathara VG. Evaluation of antioxidant and neuroprotective effect of date palm (*Phoenix dactylifera* L.) against bilateral common carotid artery occlusion in rats. *Indian J Exp Biol* 2011; 49(8): 627-33.
- [62] Pujari RR, Vyawahare NS, Thakurdesai PA. Neuroprotective and antioxidant role of *Phoenix dactylifera* in permanent bilateral common carotid occlusion in rats. *J Acute Dis* 2014; 3(2): 104-14.
- [63] Majid AS, Marzieh P, Shahriar D, Zahed SK, Pari KT. Neuroprotective effects of aqueous date fruit extract on focal cerebral ischemia in rats. *Pak J Med Sci* 2008; 24(5): 661-5.
- [64] Attah AF, O'Brien M, Koebach J, Sonibare MA, Moody JO, Smith TJ, Gruber CW. Uterine contractility of plants used to facilitate childbirth in Nigerian ethnomedicine. *J Ethnopharmacol* 2012; 143(1): 377-82.
- [65] Fallahi S, Rajaei M, Malekzadeh K, Kalantar SM. Would *Phoenix Dactylifera* Pollen (palm seed) be considered as a treatment agent against Males' infertility? A systematic review. *Electronic physician* 2015; 7(8): 1590-6.
- [66] Tahvilzadeh M, Hajimahmoodi M, Rahimi R. The role of date palm (*Phoenix dactylifera* L) pollen in fertility. *J Evid Based Complementary Altern Med* 2016; 21(4): 320-4.
- [67] Ali BH, Bashir AK, Alhadrami G. Reproductive hormonal status of rats treated with date pits. *Food Chem* 1999; 66(4): 437-41.
- [68] Ben Abdallah F, Dammak I, Mallek Z, Attia H, Hentati B, Ammar-Keskes L. Effects of date seed oil on testicular antioxidant enzymes and epididymal sperm characteristics in male mice. *Andrologia* 2009; 41(4): 229-34.
- [69] Mehraban F, Jafari M, Akbartabar Toori M, Sadeghi H, Joodi B, Mostafazade M, Sadeghi H. Effects of date palm pollen (*Phoenix dactylifera* L.) and *Astragalus ovinus* on sperm parameters and sex hormones in adult male rats. *Iran J Reprod Med* 2014; 12(10): 705-12.
- [70] El-Kashlan AM, Nooh MM, Hassan WA, Rizk SM. Therapeutic potential of date palm pollen for testicular dysfunction induced by thyroid disorders in male rats. *PLoS One* 2015; 10(10): e0139493.
- [71] Braga FC, Serra CP, Júnior NSV, Oliveira AB, Côrtes SF, Lombardi JA. Angiotensin-converting enzyme inhibition by Brazilian plants. *Fitoterapia* 2007; 78(5): 353-8.
- [72] He FJ, MacGregor GA. Beneficial effects of potassium on human health. *Physiologia plantarum* 2008; 133(4): 725-35.
- [73] Tang Z-X, Shi L-E, Aleid SM. Date fruit: chemical composition, nutritional and medicinal values, products. *J Sci Food Agric* 2013; 93(10): 2351-61.

- [74] Alsaif MA, Khan LK, Alhamdan AAH, Alorf SM, Harfi SH, Al-Othman AM, Arif Z. Effect of dates and gahwa (Arabian coffee) supplementation on lipids in hypercholesterolemic hamsters. *Int J Pharmacol* 2007; 3(2): 123-9.
- [75] Patel MD, Thompson PD. Phytosterols and vascular disease. *Atherosclerosis* 2006; 186(1): 12-9.
- [76] Ahmed S, Khan RA, Jamil S. Anti hyperlipidemic and hepatoprotective effects of native date fruit variety "Aseel" (*Phoenix dactylifera*). *Pak J Pharm Sci* 2016; 29(6): 1945-50.
- [77] Auger C, Teissedre P-L, Gérain P, Lequeux N, Bornet A, Serisier S, Besançon P, Caporiccio B, Cristol J-P, Rouanet J-M. Dietary wine phenolics catechin, quercetin, and resveratrol efficiently protect hypercholesterolemic hamsters against aortic fatty streak accumulation. *J Agric Food Chem* 2005; 53(6): 2015-21.